



**www.1stPestControl.ca**  
#4 - 7187 Honeyman St.  
Delta, British Columbia  
V4G 1E2

**Initial Treatment** \_\_\_\_\_

**Follow-Up Treatment** \_\_\_\_\_

## **HOW TO PREPARE FOR A COCKROACH TREATMENT**

**People and pets DO NOT need to vacate**

### **PRIOR TO TREATMENT:**

- ☐ **Remove items from upper and lower cupboards**
  - e.g. pots, pans, food items.
- ☐ **Remove items from drawers**
  - e.g. towels and utensils.
- ☐ **Remove items from counter tops**
  - e.g. small appliances, decorations, cookbooks.
- ☐ **Pull out fridge and stove (if possible)**
  - Our technicians will not be permitted to move your large appliances, however behind the fridge and stove are primary hiding locations for cockroaches.
- ☐ **Remove all old or ripped shelf paper**
- ☐ **Clean all areas where items have been removed**
  - Cleaning these areas is essential to a successful treatment. The goal of cleaning these areas is to eliminate any crumbs, grease and food particles that may have accumulated.
  - Don't forget to clean behind your fridge, stove, microwave and any other appliances you may have.

### **POST-TREATMENT:**

- ☐ **Replace all shelf paper**
  - loose or old shelf paper may have accumulated food debris over time and should be replaced with new shelf paper.
- ☐ **Ensure that all food is stored carefully**
  - Sealed in Tupperware or pest-proof containers.
- ☐ **Return all contents back to their original place**
- ☐ **Throw out garbage**
  - All garbage must be thrown out daily to limit potential food sources for the cockroaches.
- ☐ **Fix dripping taps or leaking pipes**

### **SAFETY INFORMATION:**

- ☐ **We will be applying:**
  - Maxforce Gel Bait (hydramethylnon 2.15% PCP 24240), Avert Dry Flowable (abamectin B1 0.05% PCP 28403), or Maxforce Impact Gel Bait (clothianidin 1.0% PCP 32275).
- ☐ **Avoid touching treated areas upon re-entry for as long as possible**
- ☐ **In an emergency, contact POISON CONTROL at 604-682-5050 or 1-800-567-8911**

### **ADDITIONAL INFORMATION:**

- ☐ **Please do not use insecticide sprays or harsh cleaners**
  - These items will render our treatment ineffective.
- ☐ **Additional activity after treatment**
  - You may experience activity for up to two (2) weeks after the initial treatment. If activity continues into the third week, you may need a follow-up treatment.

**IF THE ABOVE IS NOT COMPLETED IT MAY BE IMPOSSIBLE TO ELIMINATE THE PROBLEM**